

One sample test about a population mean with unknown population standard deviation

Some review....

We have talked about one sample proportion z-test:

Test Statistic:  $z = \frac{\hat{p} - p}{\sqrt{\frac{\hat{p}\hat{q}}{n}}}$       Confidence Interval:  $\hat{p} \pm z^* \sqrt{\frac{\hat{p}\hat{q}}{n}}$

One sample test about a population mean with known standard deviation:

Test Statistic:  $z = \frac{\bar{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$       Confidence Interval:  $\bar{x} \pm z^* \frac{\sigma}{\sqrt{n}}$

Now...we have a one sample test about a population mean with unknown standard deviation:

Test Statistic:  $t = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$       Confidence Interval:  $\bar{x} \pm t^* \frac{s}{\sqrt{n}}$

When is this used? When we are testing about a mean AND we do not know the population standard deviation.

Assumptions:

1. SRS
2. Normally distributed or  $n \geq 30$
3. Unknown population standard deviation

Example:


A college professor suspects that students at his school are getting less than 8 hours of sleep a night, on average. To test his belief, the professor asks a random sample of 30 students, "How much sleep did you get last night?" Assume a level of significance of 5% if not given. Here are the data (in hours):

9	6	8	6	8	8	6	6.5	6	7	9	4	3	4	5
10	6	11	6	3	6	6	10	7	8	4.5	9	7	7	6



**Example 2:**

At the Hawaii Pineapple Company, managers are interested in the sizes of pineapples grown in the company's fields. Last year, the mean weight of the pineapples harvested from one large field was 31 ounces. A different irrigation system was installed in this field after the growing season. Managers wonder how this change will affect the mean weight of pineapples grown in the field. To find out, they select and weigh a random sample of 50 pineapples from this year's crop. The mean size in ounces of the sample was 31.935 ounces and the sample standard deviation was 2.394. Has the mean weight changed?



**TABLE B  $t$  Distribution Critical Values**

df	TAIL PROBABILITY $P$											
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025	.001	.0005
1	1.000	1.376	1.963	3.078	6.314	12.71	15.89	31.82	63.66	127.3	318.3	636.6
2	.816	1.061	1.386	1.886	2.920	4.303	4.849	6.965	9.925	14.09	22.33	31.60
3	.765	.978	1.250	1.638	2.353	3.182	3.482	4.541	5.841	7.453	10.21	12.92
4	.741	.941	1.190	1.533	2.132	2.776	2.999	3.747	4.604	5.598	7.173	8.610
5	.727	.920	1.156	1.476	2.015	2.571	2.757	3.365	4.032	4.773	5.893	6.869
6	.718	.906	1.134	1.440	1.943	2.447	2.612	3.143	3.707	4.317	5.208	5.959
7	.711	.896	1.119	1.415	1.895	2.365	2.517	2.998	3.499	4.029	4.785	5.408
8	.706	.889	1.108	1.397	1.860	2.306	2.449	2.896	3.355	3.833	4.501	5.041
9	.703	.883	1.100	1.383	1.833	2.262	2.398	2.821	3.250	3.690	4.297	4.781
10	.700	.879	1.093	1.372	1.812	2.228	2.359	2.764	3.169	3.581	4.144	4.587
11	.697	.876	1.088	1.363	1.796	2.201	2.328	2.718	3.106	3.497	4.025	4.437
12	.695	.873	1.083	1.356	1.782	2.179	2.303	2.681	3.055	3.428	3.930	4.318
13	.694	.870	1.079	1.350	1.771	2.160	2.282	2.650	3.012	3.372	3.852	4.221
14	.692	.868	1.076	1.345	1.761	2.145	2.264	2.624	2.977	3.326	3.787	4.140
15	.691	.866	1.074	1.341	1.753	2.131	2.249	2.602	2.947	3.286	3.733	4.073
16	.690	.865	1.071	1.337	1.746	2.120	2.235	2.583	2.921	3.252	3.686	4.015
17	.689	.863	1.069	1.333	1.740	2.110	2.224	2.567	2.898	3.222	3.646	3.965
18	.688	.862	1.067	1.330	1.734	2.101	2.214	2.552	2.878	3.197	3.611	3.922
19	.688	.861	1.066	1.328	1.729	2.093	2.205	2.539	2.861	3.174	3.579	3.883
20	.687	.860	1.064	1.325	1.725	2.086	2.197	2.528	2.845	3.153	3.552	3.850
21	.686	.859	1.063	1.323	1.721	2.080	2.189	2.518	2.831	3.135	3.527	3.819
22	.686	.858	1.061	1.321	1.717	2.074	2.183	2.508	2.819	3.119	3.505	3.792
23	.685	.858	1.060	1.319	1.714	2.069	2.177	2.500	2.807	3.104	3.485	3.768
24	.685	.857	1.059	1.318	1.711	2.064	2.172	2.492	2.797	3.091	3.467	3.745
25	.684	.856	1.058	1.316	1.708	2.060	2.167	2.485	2.787	3.078	3.450	3.725
26	.684	.856	1.058	1.315	1.706	2.056	2.162	2.479	2.779	3.067	3.435	3.707
27	.684	.855	1.057	1.314	1.703	2.052	2.158	2.473	2.771	3.057	3.421	3.690
28	.683	.855	1.056	1.313	1.701	2.048	2.154	2.467	2.763	3.047	3.408	3.674
29	.683	.854	1.055	1.311	1.699	2.045	2.150	2.462	2.756	3.038	3.396	3.659
30	.683	.854	1.055	1.310	1.697	2.042	2.147	2.457	2.750	3.030	3.385	3.646
40	.681	.851	1.050	1.303	1.684	2.021	2.123	2.423	2.704	2.971	3.307	3.551
50	.679	.849	1.047	1.299	1.676	2.009	2.109	2.403	2.678	2.937	3.261	3.496
60	.679	.848	1.045	1.296	1.671	2.000	2.099	2.390	2.660	2.915	3.232	3.460
80	.678	.846	1.043	1.292	1.664	1.990	2.088	2.374	2.639	2.887	3.195	3.416
100	.677	.845	1.042	1.290	1.660	1.984	2.081	2.364	2.626	2.871	3.174	3.390
1000	.675	.842	1.037	1.282	1.646	1.962	2.056	2.330	2.581	2.813	3.098	3.300
$\infty$	.674	.841	1.036	1.282	1.645	1.960	2.054	2.326	2.576	2.807	3.091	3.291
	50%	60%	70%	80%	90%	95%	96%	98%	99%	99.5%	99.8%	99.9%
	Confidence level $C$											

What if we want a 95% confidence interval for the population mean?

You try!

Suppose we are testing a new antibacterial cream on a small cut made on the inner forearm. We know from previous research that with no medication, the mean healing time is 7.6 days. The claim we want to test is that the cream speeds healing time. We cut 250 randomly selected students (I know...sounds violent) and apply the cream to the wounds. The mean healing time is 7.5 days and the standard deviation is 0.9 days.